

Capetus

EXTRA



GROUP 3 FUNGICIDE

Safety Data Sheet for this product
scan the QR code or use the weblink:

https://links.jtcrop.net/UK_Capetusextra

Alternatively, contact your supplier.



MAPP 20675

For use as an agricultural fungicide for the control of stem-base, foliar and ear diseases in winter and spring wheat (also reduction of the mycotoxin deoxynivalenol), winter rye, winter and spring barley, winter and spring oats and for disease control in oilseed rape.

An emulsifiable concentrate formulation containing 125 g/L (12.8%) prothioconazole and 125 g/L (12.8%) tebuconazole.

UFI: 12X2-C00P-U004-W01H



Contains 125 g/L (12.8%) prothioconazole
and 125 g/L (12.8%) tebuconazole.
Also contains N,N-Dimethyl decanamide.

Warning

Causes skin irritation.
Causes serious eye irritation.
May cause an allergic skin reaction.
May cause respiratory irritation.
Suspected of damaging the unborn child.
Toxic to aquatic life with long lasting effects.

Do not get in eyes, on skin
or on clothing.

Avoid release to the
environment.

Wear protective gloves/
protective clothing/
eye protection/face protection.

IF SWALLOWED: Call a POISON
CENTRE or doctor/physician if
you feel unwell.

Collect spillage.
Dispose of contents/container
to a licensed hazardous waste
disposal contractor or collection
site except for triple rinsed empty
containers which can be disposed
of as non-hazardous waste.

Contains
2-[2-(1-chlorocyclopropyl)-2-
hydroxy-3-phenylpropyl]-2,4-
dihydro-3H-1,2,4-triazole-3-
thione. May produce an allergic
reaction..

**To avoid risks to human
health and the environment,
comply with the instructions
for use.**



PROTECT FROM FROST
SHAKE WELL BEFORE USE

Transport Information
UN 3082 ADR/RID Class: 9 Packaging Group: III

04/25

Net contents
5L e

Distributor: Crophethics Ltd
Lees Mill Lane, Huddersfield, HD7 5QE
Telephone: 01484 848885
www.jtcrop.com | info@jtcrop.com

Authorisation Holder:
JT Agro Ltd



Capetus

EXTRA



GROUP 3 FUNGICIDE

Safety Data Sheet for this product
scan the QR code or use the weblink:

https://links.jtcrop.net/UK_Capetusextra

Alternatively, contact your supplier.



MAPP 20675

For use as an agricultural fungicide for the control of stem-base, foliar and ear diseases in winter and spring wheat (also reduction of the mycotoxin deoxynivalenol), winter rye, winter and spring barley, winter and spring oats and for disease control in oilseed rape.

An emulsifiable concentrate formulation containing 125 g/L (12.8%) prothioconazole and 125 g/L (12.8%) tebuconazole.

UFI: 12X2-C00P-U004-W01H

Contains 125 g/L (12.8%) prothioconazole
and 125 g/L (12.8%) tebuconazole.
Also contains N,N-Dimethyl decanamide.



Warning

Causes skin irritation.
Causes serious eye irritation.
May cause an allergic skin reaction.
May cause respiratory irritation.
Suspected of damaging the unborn child.
Toxic to aquatic life with long lasting effects.

Do not get in eyes, on skin or on clothing.

Avoid release to the environment.

Wear protective gloves/
protective clothing/
eye protection/face protection.

IF SWALLOWED: Call a POISON CENTRE or doctor/physician if you feel unwell.

Collect spillage.
Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

Contains
2-[2-(1-chlorocyclopropyl)-2-hydroxy-3-phenylpropyl]-2,4-dihydro-3H-1,2,4-triazole-3-thione. May produce an allergic reaction..

To avoid risks to human health and the environment, comply with the instructions for use.



PROTECT FROM FROST
SHAKE WELL BEFORE USE

Transport Information

UN 3082 ADR/RID Class: 9 Packaging Group: III

04/25

Net contents
5L e

Distributor: Crophthetics Ltd
Lees Mill Lane, Huddersfield, HD7 5QE
Telephone: 01484 848885
www.jtcrop.com | info@jtcrop.com

Authorisation Holder:
JT Agro Ltd



IMPORTANT INFORMATION

FOR USE ONLY AS A PROFESSIONAL FUNGICIDE

Crops/situations	Maximum Individual dose (L product/ha)	Maximum number of treatments (per crop)	Latest time of application
Wheat, winter rye	1.0	2	End of flowering (BBCH 69)
Barley, oats	1.0	2	Beginning of flowering (BBCH 61)
Oilseed rape	1.0	2	56 days before harvest

Other specific restrictions:

A minimum interval of 21 days must be observed between applications.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

THE (COSH) CONTROL OF SUBSTANCES
HAZARDOUS TO HEALTH REGULATIONS MAY
APPLY TO THE USE OF THIS PRODUCT AT WORK.

SAFETY PRECAUTIONS

Operator Protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION when handling the concentrate.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) when applying the product.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH ANY CONTAMINATION from eyes immediately.

WASH HANDS AND EXPOSED SKIN before meals and after work.

IF YOU FEEL UNWELL, seek medical advice (show the label where possible).

Environmental Protection

DO NOT CONTAMINATE SURFACE WATERS OR DITCHES with chemical or used container.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 M of the top of the bank of a static or flowing water body, unless a Local Environment Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 M of the top of a ditch which is dry at the time of application. Aim spray away from water.

This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for three years.

Storage and Disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.

KEEP OUT OF REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER tightly closed in a safe place.

WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely.

DO NOT RE-USE CONTAINER for any purpose.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

CAPETUS EXTRA is a mixture of a triazolinthione fungicide and a triazole fungicide recommended for control of a wide range of diseases on winter and spring barley, winter and spring wheat, winter rye, winter and spring oats and oilseed rape.

PROTECT FROM FROST

CROPS

CAPETUS EXTRA may be used on all commercial varieties of winter and spring barley, winter and spring wheat, winter rye, winter and spring oats and oilseed rape.

RATE OF USE

Apply CAPETUS EXTRA at 1.0 litre per hectare
The maximum number of treatments per crop is 2.

APPLICATION

Water volume

Apply CAPETUS EXTRA in 100-400 litres water per hectare. The higher spray volumes are recommended where the crop is dense or disease pressure/risk is high to ensure good penetration to the lower leaves and stem bases. Disease control may be compromised by reducing water volumes, where good spray coverage is difficult to achieve. A spray pressure of 2-3 bar is recommended.

Spray quality

Apply as a MEDIUM spray quality (as defined by BCPC).

Latest Permitted Timing

In wheat and rye CAPETUS EXTRA may be applied at any stage up to the end of flowering (BBCH 69) and in barley and oats up to beginning of flowering. In oilseed rape CAPETUS EXTRA may be applied up to 56 days before harvest.

Mixing

Thoroughly shake the pack before use.
Add the required quantity of CAPETUS EXTRA to the half-filled spray tank with the agitation system in operation and then fill to the required level.
Continue agitation at all times during spraying and stoppages until the tank is completely empty. Spray immediately after mixing.

General

Sprayers should be thoroughly cleaned with water and detergent after use, and filters and jets checked for damage and blockages.

Boom height should be adjusted to ensure even coverage of the crop, particularly at later growth stages. The correct height is one at which the spray from alternate nozzles meets just above the crop, in dense crops, at later growth stages, higher water volumes should be used.

DISEASES CONTROLLED

Wheat

Eyespot (reduction of the incidence and severity), *Septoria* (leaf and glume blotch), powdery mildew*, yellow rust, brown rust, tan spot, ear disease complex (*Fusarium* ear blight (reduction of deoxynivalenol) and reduction of sooty moulds).

Barley

Eyespot (reduction of the incidence and severity), powdery mildew, yellow rust, brown rust, ear disease complex (*Fusarium* ear blight* and reduction of sooty moulds), *Rhynchosporium*, net blotch.

Rye

Eyespot (reduction of the incidence and severity), powdery mildew, brown rust, *Rhynchosporium*.

Oats

Eyespot, crown rust and mildew

Oilseed rape

Light leaf spot*, *Phoma* leaf spot/stem canker, and *Sclerotinia* stem rot.

*CAPETUS EXTRA will provide moderate control of these diseases.

APPLICATION TIMING

A minimum interval of 21 days must be observed between applications in all crops.

CEREALS

Eyespot (*Tapesia* spp.)

Spray in the spring at the first sign of disease, from when the leaf sheaths begin to become erect until the 2nd node is detectable (GS 30-32).

Septoria Leaf Blotch and Glume Blotch (*Septoria tritici* and *Stagonospora nodorum*)

Apply before disease is established in the crop. To protect the upper leaves and ear apply CAPETUS EXTRA at full flag leaf emergence (GS 37) up to mid-flowering (GS 65). Where disease pressure remains high application may be repeated. Applications to upper leaves where *S. tritici* symptoms are present are likely to be less effective.

CAPETUS EXTRA contains two DMI fungicides. Resistance to some DMI fungicides has been identified in *Septoria* leaf blotch (*Mycosphaerella graminicola*) which may seriously affect the performance of some products. For further advice on resistance management in DMI's contact your agronomist or specialist advisor, and visit the FRAG-UK website.

Powdery Mildew (*Blumeria graminis*)

Apply CAPETUS EXTRA at the first signs of disease. Where disease pressure remains high application may be repeated. CAPETUS EXTRA will only provide moderate control of powdery mildew in wheat

Yellow Rust (*Puccinia striiformis*)

Apply CAPETUS EXTRA at the first signs of disease. A second application may be made later if re-infection occurs. Applications made to established infections are likely to be less effective.

Brown Rust (*Puccinia recondita* and *P. hordei*)

Apply CAPETUS EXTRA at the first signs of disease. A second application may be made later if re-infection occurs. Applications made to established infections are likely to be less effective.

Crown Rust (*Puccinia coronata*)

Apply CAPETUS EXTRA at the first signs of disease. CAPETUS EXTRA controls crown rust in winter and spring oats. A second application may be made later if re-infection occurs. Applications made to established infections are likely to be less effective.

Tan Spot (*Pyrenophora tritici-repentis*)

Apply CAPETUS EXTRA at the first signs of disease in spring or early summer. Where disease pressure remains high application may be repeated.

Ear Disease Complex

Apply CAPETUS EXTRA soon after ear emergence until the end of flowering (GS 59-69) for control of *Fusarium* ear blight in wheat (or moderate control in barley) and reduction of sooty moulds. Control of ear diseases can result in cleaner, brighter ears.

Through the control of ear blight, CAPETUS EXTRA effectively reduces the level of the *Fusarium* mycotoxin deoxynivalenol (DON) in wheat grain. However, where *Fusarium* levels are high, the reduction achieved may not always be sufficient to ensure that DON levels fall below the statutory limit.

Leaf Blotch (*Rhynchosporium secalis*)

Apply CAPETUS EXTRA in spring at the first signs of disease. For severe infections a second application may be necessary later.

Net Blotch (*Pyrenophora teres*)

Apply CAPETUS EXTRA at the first signs of disease in spring/early summer. For severe infections, a second application later will give most effective control when conditions remain favourable for disease development.

OILSEED RAPE

Light Leaf Spot

Apply CAPETUS EXTRA in autumn/winter (usually late October to early December) protectively. Follow up spray(s) may be required in early spring from the onset of stem elongation, depending on disease development.

Phoma Leaf spot/Stem Canker

Apply CAPETUS EXTRA in autumn at the first sign of disease. Repeat application in late autumn/winter, if disease symptoms reoccur.

Sclerotinia stem rot (*Sclerotinia Sclerotiorum*)

Apply CAPETUS EXTRA at early to full flower

RESISTANCE STRATEGY

Repeated application of CAPETUS EXTRA alone should not be used on the same crop against a high risk pathogen such as cereal powdery mildew. Tank-mixtures or alternation with fungicides having a different mode of action have been shown to protect against the development of resistant forms of disease.

CAUTION: The possible development of disease strains resistant to CAPETUS EXTRA cannot be excluded or predicted. Where such resistant strains occur, CAPETUS EXTRA is unlikely to give satisfactory control.

COMPANY ADVISORY INFORMATION

This section is not part of the Product Label under Regulation 1107/2009 (GB/NL).

CONDITIONS OF SUPPLY

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use or the weather conditions before, during or after application, which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded, and no responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.